

DIGITAL ANO-RECTAL EXAMINATION

How to perform a DARE on yourself or your partner

STEP 1 - PREPARE

- Wash your hands with soap and warm water. Dry your hands.
- Lower or remove your underpants.
- Have water based lubricant handy.
- Have a towel or paper towels handy.
- Self/partner-DAREs can be performed with disposable gloves.

STEP 2 - POSITION

Make yourself comfortable and place yourself in one of the following positions:

- Lying on your back with one or both legs raised.
- Standing on one leg raised on a chair or stool.
- Standing with your legs spread apart while in the shower.
- If your partner is performing the DARE - bend over a table or lay on your left hand side, knees to chest.

STEP 3 - DARE

- Place water-based lubricant on a pointer finger and apply to the outer anus. Gently work your lubricated finger inside the anus.
- Insert finger till you feel the canal opening up. Up to second joint of your index finger is a good rule.
- Gently apply pressure to the wall of the anus noting lumps or differences in the surface lining of the anus.
- Repeat the process around each quarter of the anal canal, until all the surface of the anus has been examined. Alternately, you can choose to perform two sweeps of 180 degrees. You may need to switch hands.
- Be thorough! It may take up to 1 minute to perform a DARE.
- Remember to wash your hands and anus afterwards with warm soapy water.





WHAT IS A DARE?

- A DARE or Digital Ano-Rectal Examination is an easy way you can check for anal abnormalities that could lead to anal cancer.
- The aim of a DARE is to identify anal lumps when they are smaller and easier to treat. A DARE can also identify other anal abnormalities including haemorrhoids, fissures and an enlarged prostate.

Self- or partner-administered DARE is not intended to replace an anal cancer examination by your doctor. You should still have your doctor conduct an annual DARE, particularly if you are 50 years and over.

WHY SHOULD I DO A DARE?

- DAREs has been shown to be effective in detecting anal cancer.
- If an anal cancer is identified early (less than 1cm) it can be surgically removed, with near 80-100% cure rates.
- If an anal cancer tumour is identified late (i.e. greater than 1cm), it will be treated with both chemotherapy and radiotherapy, which can be uncomfortable and even invasive.



WHAT CAUSES ANAL CANCER?

- Human Papillomavirus (HPV) is the most common sexually transmitted infection (STI) worldwide. There are more than 40 types of sexually transmitted HPV.
- High-risk HPV causes cervical cancer and a large proportion of cancers of the vulva, vagina, penis, anus, mouth and throat. Chronic anal HPV infection can lead to anal cancer.
- For most people in the general community, the immune system keeps HPV under control
- People living with HIV (PLHIV) however, have difficulty keeping HPV under control and may develop chronic HPV infection.
- Anal sex with multiple partners increases the risk of HPV infection and anal cancer. However, HPV is so easily spread that men and women who never have anal sex may be at risk, because HPV can be easily transmitted from adjacent genital sites.

HOW TO I REDUCE MY RISK OF ANAL CANCER?

- Stop smoking.
- Ensure your viral load remains low and CD4 count is maintained by regularly taking your HIV meds.
- Ask your doctor for an annual DARE.
- Get any lumps, bleeding, sores, or pain in your anus checked by your doctor.
- Talk with your doctor about getting vaccinated against HPV.

AFTER A DARE

While performing the DARE if you or your partner notice any:

- lumps,
- abnormalities,
- blood,
- or discharge

Contact your doctor to discuss your findings.

IF YOU'RE CONCERNED ABOUT ANAL CANCER

- Talk to your doctor. While treatment for anal cancer is uncomfortable and can even be invasive, the sooner you act, the better the outcome.
- Talk with a peer about anal cancer. Contact a Positive Life NSW Treatments Officer, for a confidential chat. Phone (02) 9206 2177 or email contact@positivelife.org.au
- For more information on HPV related anal cancer: thebottomline.org.au

“63% of anal cancers are diagnosed late in Australia.”

MORE INFORMATION

www.positivelife.org.au
free call 1800 245 677



PositiveLifeNSW
the voice of people with HIV since 1988

Acknowledgments:

Lance Feeney, Craig Cooper, David Crawford, Dr Richard Hillman, Dr Mary Poynter, Dr Rick Varma, Dr Carmella Law, Dr Dave Templeton, Rebecca Houghton, Katy Roy, Alan G Nyitray, et al. A phase 2 clinical study to assess the feasibility of self and partner anal examinations to detect anal canal abnormalities including anal cancer, *BJM Publishing Group*, 2017

Image source: Rajab, T. K., Bordeianou, L. G., von Keudell, A., Rajab, H., & Zhou, H. (2018). Digital Rectal Examination and Anoscopy. *New England Journal of Medicine*, 378(22)

Images used are stock photos and do not imply any HIV status, sexuality or behaviours.