

# HPV & ANAL CANCER

Talk to your doctor and get checked out for anal cancer



*"HIV positive gay men have about 100 times the risk of developing anal cancer compared to the general population."*

## THE FACTS

- Human Papilloma Virus (HPV) is the most common sexually transmitted infection worldwide.
- There are more than 100 types of HPV. Around 40 types are passed on through sexual contact and can infect the cells of the anus (arse).
- About 14 of these types of HPV can cause anal cancer or anal lesions that can develop into cancer. These types are called 'high risk HPV'.
- Anal HPV infection is very common in gay men.
- Research conducted in Sydney found about three quarters of HIV positive men had high risk anal HPV.
- Why some anal HPV infections can lead to cancer but the majority of infections do not, is not currently known.
- Many pre-cancerous lesions never progress to cancer.
- Early detection means treatment is associated with greater survival rates than treatment at later stages.

## RISK FACTORS

- Gay men are five to 20 times more likely than heterosexual men to develop anal cancer.
- HIV positive gay men have about 100 times the risk of anal cancer compared with the general population.
- HIV positive women have about 15 times the risk of anal cancer compared with the general female population.
- Women who have had previous cervical dysplasia (abnormal cells) or cancer have about five times the risk of anal cancer compared with the general female population.
- Current low CD4 count, or in the past.
- Cigarette smoking.
- Increasing age.
- Receptive anal sex with multiple partners increases your risk of anal cancer. However, the virus is so easily spread that even men who never have receptive anal sex are still at increased risk.



*"The examination was slightly uncomfortable but it was worth doing it to know I didn't have cancer."*

## GET VACCINATED

- About 80% of young people in Australian schools are vaccinated against HPV.
- Ideally, get vaccinated before becoming sexually active.
- There is not much information about vaccination of people who are already infected with anal cancer HPV. The HPV vaccine could still protect you against strains you haven't already been exposed to.
- Vaccination can help protect you from reinfection.
- HPV vaccination is very effective in preventing HPV infection with anal high risk HPV to prevent the development of anal pre-cancer lesions in people who've been vaccinated.
- The vaccine is available to adults from your doctor. It is not subsidised by the Federal Government and will cost around \$450 over six months (approx. \$150 per dose).
- If you have private health insurance, some of this cost could be covered. Talk with your insurance provider about your health cover.

## GET CHECKED OUT

- Chances of survival are much higher when anal cancer is diagnosed early.
- Digital AnoRectal Examination (DARE) (examining the anus with a finger) by your doctor is recommended to screen for anal cancer.
- If further examination is required, your doctor can refer you to the Dysplasia and Anal Cancer Service (DACs) clinic.
- For peer support, contact Positive Life on (02) 9206-2177 or 1800 245 677

## REDUCE YOUR RISK

- Stop smoking.
- Ask your doctor for a digital anorectal examination each year.
- Get any lumps, bleeding, sores, or pain in your arse checked by your doctor.
- Talk with your doctor about getting vaccinated.

## MORE INFORMATION

[www.positivelife.org.au](http://www.positivelife.org.au)  
free call 1800 245 677



**PositiveLifeNSW**  
the voice of people with HIV since 1988

January 2016